

IMPLICATIONS OF THE SECONDARY ROLE OF CARBON DIOXIDE AND METHANE FORCING IN CLIMATE CHANGE: PAST, PRESENT, AND FUTURE

Willie Soon

**Harvard-Smithsonian Center for Astrophysics
Cambridge, Massachusetts**

Abstract: A review of the recent refereed literature fails to confirm quantitatively that carbon dioxide (CO₂) radiative forcing was the prime mover in the changes in temperature, ice-sheet volume, and related climatic variables in the glacial and interglacial episodes of the past 650,000 years, even under the “fast-response” framework where the convenient if artificial distinction between forcing and feedback is assumed. Atmospheric CO₂ variations generally follow changes in temperature and other climatic variables rather than preceding them. Likewise, there is no confirmation of the often-positing significant supporting role of methane (CH₄) forcing, which—despite its faster atmospheric response time—is simply too small, amounting to less than 0.2 W/m² from a change of 400 ppb. We cannot quantitatively validate the numerous qualitative suggestions that the CO₂ and CH₄ forcings that occurred in response to Milankovic orbital cycles accounted for more than half of the amplitude of the changes in the glacial/interglacial cycles of global temperature, sea level, and ice volume. Consequently, we infer that natural climatic variability—notably the persistence of insolation forcing at key seasons and geographical locations, taken with closely related thermal, hydrological, and cryospheric changes (such as the water vapor, cloud, and ice albedo feedbacks)—suffices *in se* to explain the proxy-derived, global and regional climatic and environmental phase-transitions in the paleoclimate. If so, it may be appropriate to place anthropogenic greenhouse gas emissions in context by separating their medium-term climatic impacts from those of a host of natural forcings and feedbacks that may, as in paleoclimatological times, prove equally significant. [Key words: glacial–interglacial cycles; water vapor, cloud-and-ice insulator, and albedo feedback; Milankovic orbital insolation forcing; atmospheric CO₂ and CH₄ forcing.]

INTRODUCTION

This paper addresses four issues that are fundamental to the debate concerning the role of CO₂ and CH₄ greenhouse gases in the forcing of past glacial–interglacial cycles and the role of such forcing for present and future considerations of climate change. These issues are addressed under four headings: (1) the relationship between atmospheric CO₂ and CH₄ concentrations, temperature, and ice-sheet volume; (2) atmospheric CO₂ radiative forcing as an amplifier of glacial–interglacial climate change; (3) glacial–interglacial climate change, involving a comparison of the role of orbitally moderated insolation forcing at local and regional scales with the effects of global radiative forcing from changing CO₂ concentrations; and (4) problems with CO₂ amplification of global mean temperature, especially the paucity of support provided by computer simulations and several general circulation models. These issues are addressed with reference to a wide range of recent and ongoing

research, both for and against the primary role of greenhouse gases in climate change scenarios. In conclusion, the paper stresses that changes in insolation in climatically sensitive zones exceed several-fold the global radiative forcing of climate change by CO₂ and CH₄, and that local and regional responses to insolation forcing decide the primary climatic feedbacks and changes observed from past glacial–interglacial cycles, and likely to be observed in the future.

RELATIONSHIPS BETWEEN ATMOSPHERIC CO₂ AND CH₄ CONCENTRATIONS, TEMPERATURE, AND ICE-SHEET VOLUME

One of the most notable, but somewhat surprising, consensus conclusions from ice-core drilling projects and research in Arctic and Antarctic regions (e.g., Fischer et al., 2006; Masson-Delmotte et al., 2006) is the fact that the deduced isotopic temperatures lead other climatic responses, including especially the atmospheric levels of minor greenhouse gases like CO₂ and CH₄. Fischer et al. (1999) first reported that atmospheric CO₂ concentrations increased by 80 to 100 ppm some 600±400 years *after* the warming of the last three deglaciations (or glacial terminations) in Antarctica and that relatively high CO₂ levels were sustained for thousand of years during glacial inception scenarios when Antarctic temperature had dropped significantly. Later, Monnin et al. (2001) and Caillon et al. (2003) offered clear evidence that temperature change drove atmospheric CO₂ responses during more accurately dated periods near glacial terminations I (at about 18 kyr before present, BP) and III (at about 240 kyr BP), respectively.

Stenni et al. (2001) presented evidence that the rising tendency of atmospheric CO₂ can be interrupted by abrupt events like the Antarctic Cold Reversal and related Oceanic Cold Reversal starting around 14 kyr BP, so that factors like the changes in the southern ocean, high-latitude atmospheric circulation, and dust transport can quickly trigger responses in global carbon cycling. Figure 1 offers a graphic summary of the apparent consensus on atmospheric CO₂ content being driven by change in temperature using the Vostok data first published by Petit et al. (1999). The latest 650 kyr-long record from the EPICA Dome Concordia effort, reported by Siegenthaler et al. (2005), remarkably confirmed the CO₂-temperature lag relation over terminations V, VI, and VII, covering the period from 400 to 650 kyr BP. The analyses by Delmotte et al. (2004) also confirmed that atmospheric CH₄ systematically lags the proxy for Antarctic temperatures by 1100±200 years, adopting the long time scale windows of 50 to 400 kyr for the lead-lag analyses.

Additionally, there are new and innovative analyses of both high-resolution ice and gas data from Antarctica and Greenland (Ahn and Brook, 2007; Loulergue et al., 2007; Kobashi et al., 2007) that are able to better time the complicated relation of atmospheric CO₂ and CH₄ with local and regional temperatures at both locations for specific abrupt warming and cooling events like the Dansgaard-Oeschger warming, Heinrich iceberg rafting, or even the 8.2 kyr cooling events in Greenland. Ultimately, certain local and regional climatic variations and changes must be responsible for driving the responses of the global carbon budget cycles—otherwise, some extra-terrestrial factors and/or extraordinary scenarios would need to be invoked to explain the trigger and maintenance of the atmospheric CO₂ and




Fig. 1. Vostok temperature and atmospheric CO₂ history for the past 420 kyr, from Petit et al. (1999), showing that Antarctic warming tends to lead the rise in CO₂ concentrations by several hundred years during the last four deglaciations (upper panel), and that relatively high CO₂ levels can be sustained for thousands of years during glacial inception scenarios when the temperature has dropped significantly (lower panel; see also Fischer et al., 1999). Cuffey and Vimeux (2001) and Vimeux et al. (2002) showed that the co-variation of the Vostok atmospheric CO₂ and isotopic temperature, once corrected for effects from changes in moisture source for the temperature, came to much closer timing agreement for the last 150 kyr BP, but as discussed under heading (1), the atmospheric CO₂ content must still somehow be controlled by other local and regional climatic variables.

CH₄ variations every 100 kyr or so for the past 650 kyr. Such a picture is not inconsistent with the recent comprehensive review by Peacock et al. (2006) that manages to specify a combination of distinct climatic, oceanic, and biochemical processes that could explain the co-variation of atmospheric CO₂ during a 100 kyr glacial–interglacial transition.

Synthesis studies by Shackleton (2000), Mudelsee (2001), Pepin et al. (2001), and Ruddiman and Raymo (2003) pointed out that both Antarctic temperatures and atmospheric CO₂ concentrations significantly lead the changes in the large ice sheets of the Northern Hemisphere by at least a few thousand years to as long as 14 kyr over the full 420 kyr of the Vostok record. The scenario that seems most reasonable is that external orbital insolation forcing triggered the fast and large changes in the air temperature in the Antarctic/Southern Hemisphere which, in turn, caused responses in deep-ocean properties (including changes in both the temperature and the volume of southern-sourced deep-water filling the ocean basins; Skinner, 2006) and in the global carbon cycle leading to the changing levels of atmospheric CO₂, which ultimately acted as an amplifier for glacial–interglacial ice-volume variability. Several recent pollen, glacial, and reflectance (i.e., natural gamma rays) records from the mid-latitude Southern Hemisphere clearly support such a scenario, with insolation changes in the Southern Hemisphere occurring first and then followed by related thermal, chemical, hydrologic, and cryospheric responses (Carter and Gammon, 2004; Suggate and Almond, 2005; Vandergoes et al., 2005; Sutherland et al., 2007).

Alley et al. (2002) and Johnston and Alley (2006) offered a different scenario by emphasizing a Northern Hemisphere thermal lead or trigger (see also the new evidence and discussion for the past 360 kyr in Kawamura et al., 2007), instead of a southern-lead scenario, for the consequential chains of variations in solar insolation, air and sea temperatures, CO₂, and global ice volume (mainly of the bulk continental ice sheet in the Northern Hemisphere), but this alternate picture clearly also proposes atmospheric CO₂ as the essential amplifier of the warming and cooling and hence the waxing and waning of global ice volume.

ATMOSPHERIC CO₂ RADIATIVE FORCING AS AN AMPLIFIER OF GLACIAL–INTERGLACIAL CLIMATE CHANGE

It is still unclear as to whether such a plausible CO₂-amplification scenario can be quantitatively confirmed with evidence available to date. The very long lead time by the radiative forcing of atmospheric CO₂ requires better clarification through physical modeling to account for the full dynamics of: (1) the great continental ice sheets—that is, those that can be sustained for 10 kyr in the phase of maximum ice extent, as deduced for the Laurentide Ice Sheet during the Last Glacial Maximum (LGM; Dyke et al., 2002), or for the extension of the ice sheet over the Barents and Kara Seas into the Eurasian continent (Svendsen et al., 1999); and (2) the ice conditions around the Arctic Ocean (Norgaard-Pedersen et al., 2003; Martinson and Pitmann, 2007) and their interactions with the coupled dynamics of the ocean and atmosphere, especially in the tropics (e.g., Ashkenazy and Tziperman, 2006; Roe, 2006).

It is also difficult to presume any significant amplifying role by atmospheric CO₂ for the extremely large winter cooling of 28°C over the Greenland–Arctic–North Atlantic area during the Younger Dryas event recently hypothesized by Broecker (2006). Similarly Barrows et al. (2007) noted several periods of rapid SST changes in their southern mid-latitude records around 58–38 kyr BP (i.e., Oxygen Isotope Chronozone or Marine Isotope Stage, MIS, 3) and 19.5–18.5 kyr BP (i.e., where a rapid warming of 5°C in 130 years was recorded at 19.4 kyr BP and a similarly large and rapid cooling starting around 19.2 kyr BP) that are simply difficult to explain by atmospheric CO₂ forcing.

More importantly, Roe (2006) showed that, if one adopts the rate of change of global ice volume rather than the absolute ice volume as a physically more direct measure of ice-sheet dynamics and its thermal connections to summertime temperature and local summertime insolation radiation, the sketched scenario of changing CO₂ leading the change in the global ice volume or ice sheet may not be correct. Instead, careful analyses by Roe found that atmospheric CO₂ actually lags the rate of global ice-volume change by a few thousand years, or CO₂ is at most synchronous with the rate of change in global ice volume. Roe concluded that “. . . variations in melting precede variations in CO₂. . . . This implies only a secondary role for CO₂. . . . variations [which] produce a weaker radiative forcing than the orbitally-induced changes in summertime insolation—in driving changes in global ice volume” (pp. 1 and 4). These issues will be discussed further in the following sections.

Another important empirical consensus that has recently emerged is the fact that older estimates of no more than 1–2°C changes in the tropical sea surface temperatures (SST) during the Last Glacial Maximum (LGM) around 21 kyr BP, or during other glacial–interglacial transition periods, may have been largely underestimated (e.g., Schrag et al., 1996; Peltier and Solheim, 2004). Lea et al. (2000) showed that the equatorial SSTs in the core area of the western Pacific warm pool, a region that could be a sensitive indicator of global carbon cycling, were $2.8^{\circ} \pm 0.7^{\circ}\text{C}$ colder at the LGM than at present. Visser et al. (2003) recently suggested that SST around the Indo-Pacific warm pool area by the Makasar Strait varied by about 3.5–4°C during the last two glacial–interglacial transitions. Barrows and Juggins (2005) synthesized the proxy SST records from about 165 cores for oceans around Australia, including the Indian Ocean, and their compilations showed a cooling of up to 4°C in the tropical eastern Indian Ocean and up to 7° to 9°C in higher latitude regions of the southwest Pacific Ocean during the LGM. Three new alkenone-derived SST records from the midlatitude Southern Hemisphere, presented by Pahnke and Sachs (2006), essentially confirmed the large amplitude change in SST between glacial and interglacial transitions. The consequence of these large-amplitude SST changes in the tropics and higher latitudes has led Visser et al. (2003) to suggest that a substantial portion of the 80 ppm change in atmospheric CO₂ during a glacial–interglacial transition can be simply explained by a direct change in CO₂ solubility in sea water as a function of SST change (see Peacock et al., 2006, for additional processes involving changes in sea level, oceanic circulation, and related chemical and biological responses). Peltier and Solheim (2004) offer a quantitative estimate by explaining “more than 60%” of the observed changes in glacial–interglacial CO₂ contents from air bubbles trapped in ice cores.

There are two additional roles often assigned to CO₂ radiative forcing that highlight the remarkable sensitivity of climate-system behavior and evolution to atmospheric CO₂ (e.g., Saltzman et al., 1993). We will only summarize these briefly here.

First, there are theoretical speculations, on even longer geologic timescales, about the formation of large icesheets in both hemispheres since about 2.7 million years BP (or late Pliocene transition; e.g., overviews in Droxler and Farrell, 2000; Crowley and Berner, 2001; Lisiecki and Raymo, 2007), and about the transition from the “41 kyr world” to the “100 kyr world” (i.e., actually more like ice age cycles every 80 kyr to 120 kyr or so; Liu and Herbert, 2004; Huybers and Wunsch, 2005) starting around 780 kyr ago or in the mid-Pleistocene transition, or MPT (Berger et al., 1999; Clark et al., 2006, described the MPT as a broader transitional variability zone from 1250 kyr to 700 kyr BP). It was suggested that the two phenomena/events happened because the Earth’s climate system had been undergoing gradual global cooling from a systematic decrease in atmospheric CO₂ until a dynamical air–sea–ice sheet interaction threshold was crossed. However, the evidence for an overall cooling since 780 kyr BP is not that strong, and the idea has been recently challenged by de Garidel-Thoron et al. (2005). Furthermore, non-CO₂-related explanations involving changes in basal conditions under ice sheets (Clark et al., 1999) may be significant. In addition, the MPT may simply be a dynamic system response to the continuous obliquity pacing as shown and discussed by Liu and Herbert (2004), Huybers and Wunsch (2005), and Huybers (2007). As for the explanation of the late Pliocene cooling, Ravelo et al. (2004) argued that the transition from warm early to middle Pliocene conditions to increasing glaciation in the Northern Hemisphere during the late Pliocene is linked more to nonlinear coupled dynamics of ocean and atmosphere that altered meridional heat and moisture transfers than to any persistent cooling or threshold-crossing tendency due to a decrease in atmospheric CO₂ over the past 3 million years. In two related modeling studies, Barreiro et al. (2006) and Fedorov et al. (2006) found that the physical explanation for great warmth during the early to mid-Pliocene from 5 to 3 million years BP may be connected to the reality of a permanent, rather than intermittent, El Niño condition (see, however, the challenge by Haywood et al., 2007). Such a condition would plausibly involve the collapse of trade winds along the equator, with an attendant large decrease in low-level stratus clouds, a large increase in incoming solar radiation, and an increase in atmospheric water-vapor feedback in heating up the tropical atmosphere and ocean. Huybers and Molnar (2007) recently concluded that a long-term cooling trend in the eastern tropical Pacific alone (which they clearly distinguished from explanations specifying high CO₂ in the early-to-mid-Pliocene), rather than a permanent El Niño scenario, may be sufficient for explaining increased glaciation since 3 million years BP.

The second prominent effect by CO₂ radiative forcing has been framed as follows. Two new studies have suggested that the observed variation of about 30 ppm in CO₂ concentrations during interglacial periods may have contributed significantly to the thermal and moisture instabilities that eventually drove glacial advances (Vettoretti and Peltier, 2004; Kubatzki et al., 2006). But the empirical basis for such a strong nonlinear effect of CO₂ forcing on the climate-system evolution

and change is not strong considering the small amplitude of CO₂ radiative forcing (see also comments on p. 275 in Khodri et al., 2003), the CO₂-temperature lagged-response relation, and even the actual simulated results (i.e., rather small relative differences in the near-surface global temperature of no more than 0.5°C and inland ice sheet area of no more than 0.5 million km² over North America shown in Fig. 9 of Kubatzki et al., 2006, for two different CO₂ radiative forcing scenarios). In contrast, Bauch and Kandiano (2007) point to significant differences in surface ocean conditions during the last interglacial (~130 kyr BP) and the Holocene, while invoking significant variabilities on centennial and millennial time scales during interglacials in response to intrinsic variations in solar irradiance outputs and subsequent amplification through mechanisms like solar effects on the distribution and transport of Arctic sea ice and changes in the meridional overturning circulation of the North Atlantic, as proposed earlier by Bond et al. (2001) for the Holocene.

GLACIAL-INTERGLACIAL CLIMATE CHANGE: THE ROLE OF ORBITALLY MODERATED INSOLATION FORCING AT LOCAL AND REGIONAL SCALES COMPARED WITH THE EFFECTS OF GLOBAL RADIATIVE FORCING FROM CHANGING CO₂ CONCENTRATION

This section and the one that follows focus more narrowly on the inability to find quantitative support for the putative role of CO₂ radiative forcing in the observed glacial-interglacial cycles of global ice volume and temperatures over the past 650 kyr as dictated by the latest EPICA Dome Concordia's records of climatic variability and trace-gases history (EPICA community members, 2004; Siegenthaler et al., 2005; Spahni et al., 2005).

Soon et al. (2001, p. 261) earlier cautioned against the premature rejection of the role of changes in solar radiation (i.e., from both orbital motion-induced and intrinsic solar magnetism-caused variability) in favor of the rather simplistic, and very possibly incorrect, picture of the domination of the climate system by changes in global radiative forcing related to the man-made emission of atmospheric greenhouse gases like CO₂ and CH₄. The caution was partly related to the illustrative and yet successful modeling experiments by Posmentier (1994) that showed the nonlinear dynamical responses of ice sheets to the lone modulation in seasonality induced by solar insolation.

Figure 2 shows the sharp contrast in the amplitude of variability between global annual-mean insolation and daily summer insolation at 65°N (Laskar et al., 1993). It shows the remarkably small changes of no larger than 0.6 W/m² in the net global solar radiation induced by the orbital evolution of the Earth around the Sun over the past 1 million years. This estimate, however, does not account for intrinsic changes to the Sun's irradiance as modulated by solar magnetic activity; an amplitude change of a few W/m² over a million years cannot be ruled out. But Figure 2 confirms that large changes in summer solar insolation at a key ice-nucleation location like 65°N (Roe, 2006) are clearly much larger than the relatively small global radiative forcing by changes of CO₂ for the glacial-interglacial transitions of the past 650 kyr, estimated to be 2–3 W/m². From the perspective of the climatic system, local summer insolation, as long as the persistency is guaranteed by locked-in




Fig. 2. Sensitivity of the Earth's climate to incoming radiation from the Sun over the past million years, based on calculations by Laskar et al. (1993), accounting (only) for geometric changes in the Sun–Earth orbit. First, the climate receives and reacts to local midsummer insolation (here taken at July 21 at 65°N: lower panel). The influence of persistent, daily, localized insolation at midsummer values is demonstrably of very much greater climatic effect, and hence more relevant in assessing the contribution of insolation to the paleoclimate, than the global annual mean insolation (upper panel) or the paleoclimatological forcing from CO₂ (of about 2 to 3 W/m²; Joos, 2005). (For a more direct comparison of solar insolation quantity presented here to, say, the radiative forcing from atmospheric CO₂, one needs to weigh in the additional effect of reflection of sunlight by the Earth system.)

orbital motion, is a highly relevant physical quantity for studying and quantifying the local and regional responses of the thermal and hydrologic variables, and may ultimately explain the apparent global statistics of regional weather and climate. Vandergoes et al. (2005) and Sutherland et al. (2007) have emphasized the important role of local summer solar insolation forcing rather than any global annual mean measure, within the context of Southern Hemisphere–lead scenario discussed earlier. Lorenz et al. (2006) supported the key role of local and regional

insolation changes and emphasized the nonlinear changes in the entire seasonal cycle of insolation for the spatial heterogeneity of Holocene climate trends. Empirical support for this view comes from the apparent dominance of the global radiative forcing of 2 W/m^2 (e.g., Joos, 2005) estimated from the 80 ppm change in atmospheric CO_2 over the glacial–interglacial cycles of the past 650 kyr, and yet no clear climatic response can be confidently or completely linked to CO_2 as discussed here and in the following section.

A diametrically opposite conclusion has been reached by Archer and Ganopolski (2005), arguing for the great receptivity of the climate system to global radiative forcing by CO_2 when contrasted with the orbital insolation forcing. Their appeal to CO_2 forcing modulation of the threshold of ice age cycles is not dissimilar to some of the dramatic, but qualitative, ideas reviewed earlier. This major disagreement should motivate further serious scientific inquiry. In the present brief paper concerning the quantitative role of CO_2 in glacial–interglacial changes, it is necessary to postpone a more complete synthesis of the external solar and interstellar forcing in accounting for: (1) intrinsic variability of the Sun from its thermo-nuclear and magnetic history (Gough, 1990, 2002; Turck-Chieze et al., 2005; with important insights from geological archives, e.g., Sharma, 2002; Lal et al., 2005; Bard and Frank, 2006); (2) both the long-term (Berger, 1978; Laskar et al., 2004) and shorter term (i.e., from multi-years to decades to centuries; Loutre et al., 1992) perturbations of the Earth's orbital geometry with respect to the Sun; and (3) even for any intrinsic variability related to the local interstellar (Frisch and Slavin, 2006; Muller et al., 2006; Scherer et al., 2006) and galactic (Cox and Loeb, 2007) environments.

However, most constructions of physical theory and modeling of glacial and interglacial changes (Kukla and Gavin, 2005; Roe, 2006; Tziperman et al., 2006; Huybers, 2007; Martinson and Pitman, 2007) do not require CO_2 to be a predominant forcing agent but, instead, strongly hint at both the necessary and sufficient conditions of orbital insolation forcing, its persistency, and its pacing role through nonlinear phase locking. A direct comparison of the 80 ppm change in atmospheric CO_2 for a radiative forcing of about 2 W/m^2 (e.g., Joos, 2005), with 10 W/m^2 summertime shortwave forcing, after properly folding in the albedo of melting ice and summer half-year insolation variation (Roe, 2006), provide us with a clear hint about the secondary role of CO_2 in setting the trend in climate change and other related responses during the glacial–interglacial transitions of the past 650 kyr. Neither does the estimate for radiative forcing of 0.2 W/m^2 (e.g., Joos, 2005) by atmospheric CH_4 change of about 400 ppb over the 100 kyr glacial–interglacial cycle suggest a very prominent role by CH_4 , either in isolation or in combination with CO_2 .

At this stage, it is relevant to emphasize that the popular scenario for potential episodic releases of methane hydrates, to act as a strong positive feedback commonly tied to seed atmospheric warming by CO_2 , may not be so straightforward. First, Milkov (2004) has cautiously lowered the previously accepted high estimate of global hydrate-bound gas from $21 \times 10^{15} \text{ m}^3$ of methane (or about 10,000 Gt of methane carbon) to a much lower range between 1 to $5 \times 10^{15} \text{ m}^3$ of methane (or about 500–2500 Gt of methane carbon). Next, Cannariato and Stott (2005) have recently challenged the possibly incorrect interpretation of the large $\delta^{13}\text{C}$

excursions in records of planktonic and benthic foraminifera as clathrate-derived methane release. Nor could a careful examination of the atmospheric methane carbon isotope ratio ($\delta^{13}\text{CH}_4$) from the western Greenland ice margin spanning the Younger Dryas–Preboreal transition find support for either catastrophic or gradual marine clathrate emissions (Schaefer et al., 2006). Finally, Bhaumik and Gupta (2007) have recently identified five major episodes of methane release since 3.6 million years BP in the ODP 997A site located on the crest of the Blake Outer Ridge (about 200 km off Georgia and South Carolina, USA). These they link to reduced hydrostatic pressure connected to lowered sea levels and intense glacial events, roughly coinciding with increased glaciation in the Northern Hemisphere.

Thus, many numerical attempts (shown below) to quantify the impacts from variations in two minor greenhouse gases, CO_2 and CH_4 , simply do not confirm their predominant roles in explaining the large amplitude changes in thermal, hydrologic, and cryospheric history during glacial–interglacial transitions. This failure to link quantitatively the seed CO_2 -induced thermal perturbations to large hydrologic and cryospheric responses is the necessary reason for questioning the CO_2 -amplifier idea. Persistent solar insolation forcing at key seasons and geographical locations, and closely related thermal, hydrological, and cryospheric changes (including the water-vapor, cloud, and ice–albedo feedbacks), may be sufficient to explain regional and global climatic changes during glacial–interglacial transitions.

PROBLEMS WITH CO_2 AMPLIFICATION OF GLOBAL MEAN TEMPERATURE: COMPUTER SIMULATIONS AND QUANTITATIVE DATA

Why is the climatic role of CO_2 radiative forcing deemed so hard to confirm? Figure 3 may help explain the inherent difficulty in confirming any radiative impact from added CO_2 forcing using the deduced global net longwave (LW) fluxes available from the International Satellite Cloud Climatology Project (ISCCP) over the 18-year span from July 1983 through June 2001, despite some known data limitations for ISCCP (e.g., Kato et al., 2006; Evan et al., 2007). Over that period, the CO_2 increase is estimated to produce an equivalent global LW forcing of only about 0.3 W/m^2 and this amount is practically not discernible from the large interannual variability of LW fluxes at either the surface, the atmospheric air column, or even the top of the atmosphere. It is understood in scientific discussions, but popularly least appreciated, that the argument for a significant role from added radiative forcing by anthropogenic emissions of CO_2 rests on the assumption that over a sufficiently long interval, say over several decades to a century, the Earth's climate system will be in some form of "equilibrium" state where all the intrinsic LW flux fluctuations shown in Figure 3 will cancel almost exactly to zero. The cancellation would in turn allow the detection and hence all related climatic manifestations of, for example, a systematic increase of about 4 W/m^2 of net radiative LW forcing (3.5 to 4.2 W/m^2 in the CO_2 forcing parameterization of 20 GCMs examined by Forster and Taylor, 2006) from the doubling of CO_2 content roughly over 70 years' time (i.e., a compounded rate of CO_2 increase at 1% per year).

It is thus widely accepted that, although atmospheric CO_2 and CH_4 contents during the glacial–interglacial cycle are a response to climate-induced perturbations to




Fig. 3. Global net longwave (LW) fluxes deduced for the surface, atmospheric column, and the top of the atmosphere from July 1983 through June 2001 in comparison to the estimated radiative forcing of 0.3 W/m^2 from increased anthropogenic CO_2 over the same 18-year time span, as well as the 4 W/m^2 estimate for a doubling of atmospheric CO_2 , which roughly extends over a 70-year period if one compounded the CO_2 concentration increase at a rate of 1% per year (adapted from the original figure shown in International Satellite Cloud Climatology Project, ISCCP, web page <http://isccp.giss.nasa.gov/projects/flux.html>, with technical discussion in Zhang et al., 2004). Recent and projected CO_2 forcing are likely to be confused by the LW variations produced by internal variability and solar radiation-induced forcing and feedback through water vapor and cloud variations.

the global carbon and methane reservoirs, both on land and in the surface ocean, continental margin, and deep sea, the CO_2 and CH_4 radiative forcing can in turn act to strongly amplify and synchronize climatic changes across all weather regimes and climatic zones from south to north poles, ultimately producing a global warming or cooling. However, a closer look reveals that most of the claims, even in many scientific publications (i.e., from Genthon et al., 1987; Lorius et al., 1990; through Hansen et al., 2007), have not offered reliable quantitative support for the claim. The necessity of *a priori* forward calculations for the difficult task of quantifying the climatic role of CO_2 radiative forcing must be contrasted with multivariable regression analyses as performed in those cited studies, adopting rather selective variables that can easily be confused by the concepts of forcing and feedback (see below). This may be the reason for the early caution issued by Genthon et al. (1987, p. 415) that “ CO_2 changes might just be a consequence of climatic change without much effect on climatic change itself.”

Furthermore, there are clearly too many adjustable values in the estimates of radiative forcing by various other factors. For example, aerosol-dust forcing during the LGM was estimated to be $-1.0 \pm 0.5 \text{ W/m}^2$ by Hansen et al. (1993) and then later modified to $-0.5 \pm 1.0 \text{ W/m}^2$ in Hansen et al. (1997; based on the modeling study of Overpeck et al., 1996, that has since been questioned by Claquin et al., 2003). A significantly larger estimate by Claquin et al. (2003) gave a global dust-forcing value during LGM that ranges from -1.0 to -3.2 W/m^2 , with forcing at higher latitudes (poleward of 45°) from -0.9 to $+0.2 \text{ W/m}^2$ and 15°N – 15°S tropical forcing ranging from -2.2 to -3.2 W/m^2 .

It is puzzling that well-accepted roles of atmospheric water vapor and cloud feedbacks (despite the fact that both variations in the isotopic proxies δD_{ice} and $\delta^{18}\text{O}_{\text{ice}}$ from ice cores are essentially markers of large hydrologic changes) are not often factored in or discussed more seriously when CO_2 as the amplifier of glacial–interglacial warming or cooling is being considered. This is especially so because varying levels of atmospheric CO_2 largely seem to be a climatic feedback response, rather than any external forcing as envisioned in the scenario of anthropogenic emissions of CO_2 .¹ Other potentially powerful hydrologic feedbacks, like the LGM hydrologic cycle weakened by a significant lowering of the mean residence time of water vapor from excessive dust loading in the atmosphere proposed by Yung et al. (1996), also have not gained much attention compared to those from added radiative forcing by CO_2 . Paleoclimatic studies discussing the modulation of greenhouse effects by water vapor and cloud formation, e.g., over warm ocean areas and follow-up effects by orbital forcing (e.g., Gupta et al., 1996) should be a priority. Priem (1997) even went as far as to suggest that, in an early Earth comprising mainly oceans, the powerful greenhouse effects of water vapor, rather than CO_2 , could come a long way toward resolving the “faint young Sun paradox.”

Another example of potentially important feedback concerns how the seasonal cycle of surface-penetrating solar radiation is coupled to oceanic biota and the related biogeochemical emissions and atmospheric responses, especially through the production of marine biogenic dimethylsulfide (e.g., Shell et al., 2003; Vallina and Simo, 2007). Finally, beneficial insights may also be gained from studying how orbital forcing affects evolution of water and CO_2 cycles and climate on Mars (e.g., Richardson and Mischna, 2005).

The immediate question then is whether one can find a clear and dominant climatic impact signal by CO_2 radiative forcing in the glacial–interglacial transition from the current state-of-the-art modeling results from various General Circulation Models (GCMs). Let us start by imposing a CO_2 radiative forcing estimate of about 2.5 W/m^2 from the 80 ppm change, which will trigger a warming of 2 – 3°C (taking a high value of climate sensitivity of 1°C per 1 W/m^2), and that falls far short of the calibrated 10 – 12°C in Antarctic temperature change. One should note that the climate sensitivity value adopted for this simple estimate has been generous, considering the “black-body” or “nofeedback” sensitivity value of 0.3°C per W/m^2 or the

¹Even in this scenario of CO_2 emissions from using fossil fuels, the actual amount of CO_2 ultimately retained in the atmosphere is still limited by climatic, chemical, and biological factors of the global carbon cycle.




Fig. 4. The successful simulations of a significant snow accumulation in the glacially sensitive location (70°N, 80°W) around the Laurentide ice sheet area for the glacial inception scenario, with orbital forcing conditions around 115 kyr BP (compared to the present day orbital forcing case, PD) and taking into account the coupled ocean-atmosphere feedbacks, but with no change in CO₂ forcing (set at about 270 ppm) between 115 kyr BP and PD (adapted from Khodri et al., 2001). The Laurentide ice sheet was probably formed without much help from CO₂ forcing.

accepted range of values from 0.4 to 1.2°C per W/m² after accounting for net gain from all the positive and negative feedback processes (Joshi et al., 2003; Forster and Taylor, 2006). A more concrete estimate comes from recent simulations of LGM by Schneider von Deimling et al. (2006a), where the CO₂ forcing contributes about 1.8°C (or about 31%) of the total 5.7°C cooling over the globe (see also Fig. 4B in Schneider von Deimling et al., 2006b). The amount is impressive but not predominantly large. Also it is clear from the spatial pattern of change that the effects from the ice-sheet forcing (represented as an albedo effect) are clearly more extensive and variable than effects from changes in CO₂ forcing.

More importantly, the notion of what is forcing and what is feedback is sufficiently confused here and has led Hansen et al. (2007) to ponder that:

[c]limate sensitivity when surface properties are free to change . . . reveals Antarctic temperature increase of 3°C per W/m². Global temperature change is about half that in Antarctica, so this equilibrium global climate sensitivity is 1.5°C per W/m², double the fast-feedback (Charney) sensitivity. Is this 1.5°C per W/m² sensitivity, rather than 0.75°C per W/m², relevant to human-made forcings? (pp. 1944 and 1946)

The issue may not be fully resolvable for now, but it is clear that this potential double counting of radiative “forcing” effects by CO₂ would, in Hansen et al.’s

(2007) view, stand as authoritative if no contest or discussion to this problematic proposal is forthcoming. In any case, if the impacts by CO₂ radiative forcing were to be real for large glacial–interglacial transitions, one should be able to verify the large stratospheric warming by up to 7°C at 60 km predicted during the LGM by, for example, Crutzen and Bruhl (1993). It would be also important to see if the CO₂ forcing theory can explain the regional warming around the Seas of Japan and Okhotsk during the LGM (Ishiwatari et al., 2001; Seki et al., 2004) where it was suspected that the anomalously warm sea surface temperature may represent the local equilibrium of thermal energy from trapped solar radiation in shallow water under the highly stratified upper ocean condition of the LGM when the Japan and Okhotsk seas were rather isolated from the open ocean as a result of the lowered sea level. Similarly, a correct CO₂ global radiative forcing theory should also be able to account for the ice-free conditions during the LGM for the coastal oasis regions of the Bunger Hills (Gore et al., 2001) and Larsemann Hills (Hodgson et al., 2006) around East Antarctica.

Another way to assess the role of CO₂ forcing in glacial–interglacial climate change would be to study the quantitative results from variations induced by orbital forcing alone in order to find out if there is any need to invoke CO₂ as a forcing input. Figure 4 shows the successful simulation of a significant snow accumulation in a glacially sensitive location (70°N, 80°W) around the Laurentide ice sheet area for the glacial inception scenario at orbital forcing condition around 115 kyr BP (compared to the present-day orbital forcing case, PD) taking into account the coupled ocean–atmosphere feedbacks but with no change in radiative forcing by CO₂ (set at about 270 ppm) between 115 kyr BP and PD by Khodri et al. (2001). In other words, the Khodri et al. (2001) study confirms the important role of seasonality and the correct accounting of complex feedback mechanisms, involving atmospheric winds, ocean dynamics, and hydrologic cycles with little hint for the need of CO₂ forcing. These results are consistent with modeling experiments of Vettoretti and Peltier (2004). Recent glacial inception modeling experiments by Risebrobakken et al. (2007) have also essentially supported the scenario by Khodri et al. (2001), while stressing dynamics from an enhanced, rather than weakened, Atlantic meridional overturning circulation in creating a strong land–sea thermal gradient together with a strong wintertime latitudinal insolation gradient to promote increased storminess and moisture transport that feeds into the formation of the Northern European ice sheet.

Loutre and Berger (2000) further emphasized the key role of orbital solar radiation forcing in generating the glacial–interglacial cycles of ice-volume changes. The authors showed that, if time-varying CO₂ forcing is prescribed alone, their model is able to generate glacial–interglacial temperature changes, but unable to simulate the simultaneously varying ice-volume cycles of the ice ages and interglacial warm stages. In contrast, the glacial–interglacial ice-volume cycles can be generated by accounting for the orbital forcing alone with a constant level of atmospheric CO₂ lower than 220 ppm. As noted earlier, such a great sensitivity of large continental ice sheet formation to the threshold crossing at a particular low CO₂ level requires more in-depth scientific research, but one can offer a counter-example from other existing CO₂-climate modeling experiments.




Fig. 5. Perennial snow accumulation rate (in m/kyr) for low tilt angle ($\epsilon = 22.0^\circ$) and highly eccentric ($e \sin(\omega) = 0.04$ with precession parameter $\omega = 90^\circ$) orbital (i.e., near the glacial inception phase) scenarios for CO_2 levels set at 290 ppm (left panel; roughly corresponds to the termination of marine isotope stage 5; 115 kyr BP) and 260 ppm (right panel; roughly corresponds to the termination of marine isotope stage 7; 220 kyr BP); adapted from Vettoretti and Peltier, 2004). Greater CO_2 forcing yields larger snow accumulation over the Arctic Ocean (see discussion in the text).

Figure 5 shows the curious example of a more extensive snow accumulation over the Arctic sea area for a case of high CO_2 level of 290 ppm in contrast to the lower CO_2 level case of 260 ppm shown in Vettoretti and Peltier (2004). Both simulations were set with exactly the same orbital configuration of low tilt angle and high eccentricity (to emulate glacial inceptions near the terminations of MIS stages 5 and 7, respectively), so the results strictly represent the consequences of having differing levels of CO_2 radiative forcing. The results show more extensive snow accumulation, rather than less, with higher CO_2 forcing, although the rates of snow accumulation in the Canadian Arctic and coastal eastern Siberia are higher for the low CO_2 case of 260 ppm. The examination of the simulated land surface temperatures and precipitation-minus-evaporation (P-E) anomalies in the polar region shows that, although polar surface land temperatures may be warmer with less extensive cool-summer areas (i.e., latitudes with temperatures from -4 to -12°C ; see Figs. 8f and 8g in Vettoretti and Peltier, 2004) in the 290 ppm case, the positive P-E regions were slightly larger and enhanced in higher polar latitudes for the experiment with 290 ppm of CO_2 (see Figs. 9f and 9g in Vettoretti and Peltier, 2004). The results in Figure 5 may be consistent with the cryospheric moisture pump scenario for glacial inceptions being more effective at a higher CO_2 scenario, studied earlier by the same authors (Vettoretti and Peltier, 2003), in which an initial cooling at high latitudes by orbital insolation forcing caused evaporation to drop locally more quickly than precipitation, which then created a condition favoring more moisture transport into polar regions via increased baroclinic activity at mid- to high latitudes in the Northern Hemisphere summer season. The results of Vettoretti and Peltier (2004) are consistent with those of Khodri et al. (2001).

What about evidence for a greater role of CO_2 radiative forcing during deglaciation scenarios? The recent hypothesis of Martinson and Pitman (2007) does not




Fig. 6. Simulations of land-ice thickness (in m) for orbital deglaciation scenarios between LGM (21 kyr BP) and the early Holocene (11 kyr BP) with the same CO₂ level at 200 ppm (left panel → top right panel) and with the CO₂ level increased from 200 ppm at LGM to 280 ppm in the early Holocene (left panel → bottom right panel; adapted from Yoshimori et al., 2001). Orbitorally induced solar forcing causes greater land-ice response than even a significant increase in atmospheric CO₂ concentration by 80 ppm.

specify any prominent role for atmospheric CO₂ but instead details the sufficiency of a sequence of events during the last glacial termination involving the crucial role of the southward expansion of the North American and Eurasian ice sheets, sea ice cover in the North Pacific and North Atlantic, and balancing acts among Arctic freshwater budgets, salinity-driven formation of polynyas (i.e., ice-free areas), and deepwater formation in the Arctic Ocean and its subsequent overflow into the North Atlantic, as well as the incursion of warm surface water from the North Atlantic to the Arctic Ocean. This hypothesis is consistent with the dynamic changes of land and sea ice contributions to the sedimentary record from the central Arctic Ocean (Spielhagen et al., 1997) that did not invoke any contribution from CO₂ radiative forcing.

Figure 6 shows the land ice thickness simulations of the deglaciation scenario from Yoshimori et al. (2001) for orbital conditions from 21 kyr (LGM) to 11 kyr BP (early Holocene), first with a constant CO₂ at 200 ppm, and then with the level of CO₂ changed by 80 ppm for the early Holocene in order to be consistent with ice-core air bubble results. The results clearly suggest a minimal impact of added radiative forcing of CO₂ on the thickness of ice sheets on land. Yoshimori et al. did

argue for a “powerful” feedback role by CO₂ forcing in explaining glacial termination, but the authors pointed out that the effect of increasing atmospheric CO₂ from 200 to 280 ppm in their simulation leads to a nominal impact on winter air temperatures over continents adjacent to the North Atlantic. That CO₂ impact in turn contributes to ice-sheet nourishment through slightly enhanced winter precipitation, so that CO₂ acts as negative, rather than positive, feedback for ice-sheet retreat during deglaciation.

The examples in Figures 4, 5, and 6 serve only as the sufficient but not necessary condition of orbital insolation forcing in accounting almost fully for conditions and changes during the glacial–interglacial transition, without the need to invoke the argument for CO₂ as the predominant amplifier of those changes.

One cannot totally discount other contemporary studies (Weaver et al., 1998; Pepin et al., 2001; Lea, 2004) that suggested a “dominant” contribution by CO₂ radiative forcing to the observed glacial–interglacial temperature change while perhaps ignoring the large changes in global ice volume and its effects. Lea (2004; citing Hewitt and Mitchell, 1997; Weaver et al., 1998; and others) claimed that “modeling results for the glacial oceans support the hypothesis that CO₂ variations are the dominant source of radiative forcing in the tropical ocean regions” (p. 2170). However, simulation results from Hewitt and Mitchell (1997) estimated the lowering of CO₂ at 21 kyr BP at the LGM to be a cooling of 1.4°C or about one-third of the total simulated cooling. Weaver et al. (1998) suggested that “the most important [more so than ice-sheet albedo feedbacks] of these forcings in our model is the change in atmospheric CO₂” but concurrently admitted to underestimating the “ice albedo” effects. More importantly, the model of Weaver et al. suggests that temperatures in the tropics were 2.2°C less than today and their results show apparent insensitivity to changes in oceanic circulation. These results are not consistent with larger amplitude change in tropical SST of 3.5°–4°C during the last two glacial–interglacial transitions, as deduced by Visser et al. (2003), and with the importance of oceanic feedbacks for glacial inception scenarios identified with the coupled ocean-atmosphere GCM as discussed by Khodri et al. (2001) and found in climate sensitivity experiments conducted by Vettoretti and Peltier (2004).

CONCLUSIONS

There is no quantitative evidence that varying levels of minor greenhouse gases like CO₂ and CH₄ have accounted for even as much as half of the reconstructed glacial–interglacial temperature changes or, more importantly, for the large variations in global ice volume on both land and sea over the past 650 kyr. This paper shows that changes in solar insolation at climatically sensitive latitudes and zones exceed the global radiative forcings of CO₂ and CH₄ by severalfold, and that regional responses to solar insolation forcing will decide the primary climatic feedbacks and changes (see also independent research and conclusions by Kukla and Gavin, 2005; Lorenz et al., 2006; Roe, 2006).

Persistent orbitally moderated insolation forcing is, therefore, likely to be the principal driver of water-vapor cycling, and the cloud-and-ice insulator and albedo feedbacks. Such a forcing-response scenario has not received enough attention in

current research (but with notable exceptions, e.g., Dong and Valdes, 1995; Gupta et al., 1996; Broecker, 1997; Greene et al., 2002; Leduc et al., 2007). A host of other forcings and feedbacks, including dust-and-aerosol forcings, oceanic circulation, and vegetation-cover feedbacks, have not been soundly quantified. The forcing from intrinsic variation of solar radiation and magnetic activity has also been almost entirely ignored in this paper, but several recent studies are beginning to document and formulate testable climatic responses on multidecadal to centennial to millennial timescales resulting from this particularly complex expression of solar change (e.g., Bond et al., 2001; Holzkamper et al., 2004; Mayewski et al., 2004; Holzhauser et al., 2005; Maasch et al., 2005; Soon, 2005; Scherer et al., 2006). There are still questions about how orbital forcings explain glaciation and deglaciation over the past few million years (Roe, 2006; Tziperman et al., 2006; Huybers, 2007; Lisiecki and Raymo, 2007), with the 100-kyr glacial–interglacial cycles not fully explained (but most likely, nonlinearly related to obliquity forcing, thereby emphasizing the key role of the insolation gradient as the driver for climatic processes and feedbacks, as discussed by Raymo and Nisancioglu, 2003; Liu and Herbert, 2004; Loutre et al., 2004; Huybers and Wunsch, 2005; Huybers, 2007).

However, the popular notion of CO₂ and CH₄ radiative forcing as the predominant amplifier of glacial–interglacial phase transitions cannot be confirmed. In this context, the graph of “radiative perturbation” during the last glacial maximum, shown as the top left panel of Figure 6.5 on p. 451 of the IPCC (2007) Working Group I report, may be gravely misleading. It suggests that *global annual mean radiative influences* by orbitally moderated insolation forcing are negligible when compared to *radiative influences* of CO₂, CH₄ + N₂O, mineral dust, continental ice and sea level, and vegetation. In our opinion, the listed influences are very likely to be the responses from the initial orbital insolation forcing and its persistent effects. Provided that the deduced amplitude of 80 ppm and 400 ppb for CO₂ and CH₄ from air-bubble records is not severely underestimated, enhanced greenhouse effects from these two minor greenhouse gases cannot explain the greater part of the large climatic swings and substantial hydrologic and cryospheric changes reconstructed for the glacial–interglacial transitions over the last 650 kyr.

Our basic hypothesis is that long-term climate change is driven by insolation changes, from both orbital variations and intrinsic solar magnetic and luminosity variations. This implies natural warming and cooling variations on decades through millennia (e.g., Bond et al., 2001; Holzkamper et al., 2004; Holzhauser et al., 2005; Maasch et al., 2005; Soon, 2005), together with an eventual cooling of the Earth and an increase in ice-mass accumulation within the past and future horizons of the Holocene and the next few thousand years or so (see Kukla and Gavin, 2005, and Bauch and Kandiano, 2007, for the pioneering research and more in-depth discussion). Such a retrodiction appears consistent with proxy evidence indicating both systematic and significant cooling trends during the Holocene² in Greenland

²Our brief discussion on the Holocene changes is not intended to be complete; see additional discussion in Mayewski et al. (2004) and Lorenz et al. (2006), as well as the Holocene SST database offered by GHOST (Global Holocene Spatial and Temporal Climate Variability) at <http://www.pangaea.de/Projects/GHOST>.

(Johnsen et al., 2001), western Arctic (Kaufman et al., 2004), Nordic Seas (Andersen et al., 2004), other regions around the northeast Atlantic and Mediterranean (Marchal et al., 2002; Kim et al., 2007; Magny et al., 2007), northwest Atlantic (Sachs, 2007), northwest Africa and Gulf of Guinea (Kim et al., 2007; Weldeab et al., 2007), western tropical Pacific ocean (Stott et al., 2004), southern mid-latitude seas south of the Indian Ocean and around the Australian–New Zealand region (Ikehara et al., 1997; Pahnke and Sachs, 2006; Barrows et al., 2007), and even the Antarctic Peninsula and both coastal and inland regions of East Antarctica (Masson et al., 2000; Masson-Delmotte et al., 2004; Hodgson et al., 2006; Smith et al., 2007). This predictable tendency and currently observed reality led Sachs (2007) to the conclusion, with which we agree, that the Holocene, often considered a time of climate stability, has been characterized by large secular changes throughout the climate system. Again, following Sachs, perhaps the cooling of northwest Atlantic slope waters during the Holocene is a harbinger of climate deterioration preceding the next glaciation.

Acknowledgments: I thank Guido Vettoretti and Thomas Schneider von Deimling for professional courtesy in answering some questions concerning their papers, Masakazu Yoshimori for sharing his figures and Ph.D. thesis, and especially Eugene Avrett and Christopher Monckton for significant improvements in several drafts. I further thank the three referees, especially referee C, for constructive reviews. Than, Lien, and Julia Pham, and Benjamin and Franklin Soon are acknowledged for loving support and motivation.

REFERENCES

- Ahn, J. and Brook, E. J. (2007) Atmospheric CO₂ and climate from 65 to 30 ka B.P. *Geophysical Research Letters*, Vol. 34, L10703, doi:10.1029/2007GL029551.
- Alley, R. B., Brook, E. J., and Anandakrishnan, S. (2002) A northern lead in the orbital band: North–south phasing of Ice-Age events. *Quaternary Science Reviews*, Vol. 21, 431–441.
- Andersen, C., Koc, N., Jennings, A., and Andrews, J. T. (2004) Nonuniform response of the major surface currents in the Nordic Seas to insolation forcing: Implications for the Holocene climate variability. *Paleocoenography*, Vol. 19, PA2003, doi:10.1029/2002PA000873.
- Archer, D. and Ganopolski, A. (2005) A movable trigger: Fossil fuel CO₂ and the onset of the next glaciation. *Geochemistry Geophysics Geosystems*, Vol. 6, Q05003, doi:10.1029/2004GC000891.
- Ashkenazy, Y. and Tziperman, E. (2006) Scenarios regarding the lead of equatorial sea surface temperature over global ice volume. *Paleocoenography*, Vol. 21, PA2006, doi:10.1029/2005PA001232.
- Bard, E. and Frank, M. (2006) Climate change and solar variability: What's new under the sun? *Earth and Planetary Science Letters*, Vol. 248, 1–14.
- Barreiro, M., Philander, G., Pacanowski, R., and Fedorov, A. (2006) Simulations of warm tropical conditions with application to Pliocene atmospheres. *Climate Dynamics*, Vol. 26, 349–365.

- Barrows, T. T. and Juggins, S. (2005) Sea-surface temperatures around the Australian margin and Indian Ocean during the last glacial maximum. *Quaternary Science Reviews*, Vol. 24, 1017–1047.
- Barrows, T. T., Juggins, S., de Deckker, P., Calvo, E., and Pelejero, C. (2007) Long-term sea surface temperature and climate change in the Australian–New Zealand region. *Paleocenography*, Vol. 22, PA2215, doi:10.1029/2006PA001328.
- Bauch, H. A. and Kandiano, E. S. (2007) Evidence for early warming and cooling in North Atlantic surface waters during the last interglacial. *Paleoceanography*, Vol. 22, PA1201, doi:10.1029/2005PA001252.
- Berger, A. (1978) Long term variations of daily insulations and Quaternary climatic changes. *Journal of the Atmospheric Sciences*, Vol. 35, 2362–2367.
- Berger, A., Li, X. S., and Loutre, M. F. (1999) Modelling Northern Hemisphere ice volume over the last 3 Ma. *Quaternary Science Reviews*, Vol. 18, 1–11.
- Bhaumik, A. K. and Gupta, A. K. (2007) Evidence of methane release from Blake Ridge ODP Hole 997A during the Plio-Pleistocene: Benthic foraminifer fauna and total organic carbon. *Current Science*, Vol. 92, 192–199.
- Bond, G., Kromer, B., Beer, J., Muscheler, R., Evans, M. N., Showers, W., Hoffmann, S., Lotti-Bond, R., Hajdas, I., and Bonani, G. (2001) Persistent solar influence on North Atlantic climate during the Holocene. *Science*, Vol. 294, 2130–2136.
- Broecker, W. S. (1997) Mountain glaciers: Recorders of atmospheric water vapor content? *Global Biogeochemical Cycles*, Vol. 11, 589–597.
- Broecker, W. S. (2006) Abrupt climate change revisited. *Global and Planetary Change*, Vol. 54, 211–215.
- Caillon, N., Severinghaus, J. P., Jouzel, J., Barnola, J.-M., Kang, J., and Lipenkov, V. Y. (2003) Timing of atmospheric CO₂ and Antarctic temperature changes across termination III. *Science*, Vol. 299, 1728–1731.
- Cannariato, K. G. and Stott, L. D. (2005) Evidence against clathrate-derived methane release to Santa Barbara Basin surface waters? *Geochemistry Geophysics Geosystems*, Vol. 5, Q05007, doi:10.1029/2003GC000600.
- Carter, R. M. and Gammon, P. (2004) New Zealand maritime glaciation: Millennial-scale southern climate change since 3.9 Ma. *Science*, Vol. 304, 1659–1662.
- Claquin, T., Roelandt, C., Kohfeld, K. E., Harrison, S. P., Tegen, I., Prentice, I. C., Balkanski, Y., Bergametti, G., Hansson, M., Mahowald, N., Rodhe, H., and Schulz, M. (2003) Radiative forcing of climate by ice-age atmospheric dust. *Climate Dynamics*, Vol. 20, 193–202.
- Clark, P. U., Alley, R. B., and Pollard, D. (1999) Northern Hemisphere ice-sheet influences on global climate change. *Science*, Vol. 286, 1104–1111.
- Clark, P. U., Archer, D., Pollard, D., Blum, J. D., Rial, J. A., Brovkin, V., Mix, A. C., Pisias, N. G., and Roy, M. (2006) The middle Pleistocene transition: Characteristics, mechanisms, and implications for long-term changes in atmospheric pCO₂. *Quaternary Science Reviews*, Vol. 25, 3150–3184.
- Cox, T. J. and Loeb, A. (2007) The collision between the Milky Way and Andromeda. *Monthly Notices of the Royal Astronomical Society*, submitted (astro-ph paper arXiv:0705.1170v1).
- Crowley, T. J. and Berner, R. A. (2001) CO₂ and climate change. *Science*, Vol. 292, 870–872.

- Crutzen, P. J. and Bruhl, C. (1993) A model study of atmospheric temperatures and the concentrations of ozone, hydroxyl, and some other photochemically active gases during the glacial, the pre-industrial Holocene and the present. *Geophysical Research Letters*, Vol. 20, 1047–1050.
- Cuffey, K. M. and Vimeux, F. (2001) Covariation of carbon dioxide and temperature from the Vostok ice core after deuterium-excess correction. *Nature*, Vol. 412, 523–527.
- de Garidel-Thoron, T., Rosenthal, T., Bassinot, F., and Beaufort, L. (2005) Stable sea surface temperatures in the western Pacific warm pool over the past 1.75 million years. *Nature*, Vol. 433, 294–298.
- Delmotte, M., Chappellaz, J., Brook, E., Yiou, P., Barnola, J. M., Goujon, C., Raynaud, D., and Lipenkov, V. I. (2004) Atmospheric methane during the last four glacial–interglacial cycles: Rapid changes and their link with Antarctic temperature. *Journal of Geophysical Research*, Vol. 109, D12104, doi:10.1029/2003JD004417.
- Dong, B. and Valdes, P. J. (1995) Sensitivity studies of Northern Hemisphere glaciation using an atmospheric general circulation model. *Journal of Climate*, Vol. 8, 2471–2496.
- Droxler, A. W. and Farrell, J. W. (2000) Marine Isotope Stage 11 (MIS 11): New insights for a warm future. *Quaternary Science Reviews*, Vol. 24, 1–5.
- Dyke, A. S., Andrews, J. T., Clark, P. U., England, J. H., Miller, G. H., Shaw, J., and Veillette, J. J. (2002) The Laurentide and Innuitian ice sheets during the Last Glacial Maximum. *Quaternary Science Reviews*, Vol. 21, 9–31.
- EPICA community members. (2004) Eight glacial cycles from an Antarctic ice core. *Nature*, Vol. 429, 623–628.
- Evan, A. T., Heidinger, A. K., and Vimont, D. J. (2007) Arguments against a physical long-term trend in global ISCCP cloud amounts. *Geophysical Research Letters*, Vol. 34, L04701, doi:10.1029/2006GL028083.
- Fedorov, A. V., Dekens, P. S., McCarthy, M., Ravelo, A. C., deMenocal, P. B., Barreiro, M., Pacanowski, R. C., and Philander, S. G. (2006) The Pliocene paradox (mechanisms for a permanent El Niño). *Science*, Vol. 312, 1485–1489.
- Fischer, H., Kull, C., and Kiefer, T. (2006) Ice core science. *PAGES News*, Vol. 14, No. 1, 1–44.
- Fischer, H., Wahlen, M., Smith, J., Mastroianni, D., and Deck, B. (1999) Ice core records of atmospheric CO₂ around the last three glacial terminations. *Science*, Vol. 283, 1712–1714.
- Forster, P. M. D. and Taylor, K. E. (2006) Climate forcings and climate sensitivities diagnosed from coupled climate model integrations. *Journal of Climate*, Vol. 19, 6181–6194.
- Frisch, P. C. and Slavin, J. D. (2006) Short-term variations in the galactic environment of the Sun. In P. C. Frisch, ed., *Solar Journey: The Significance of Our Galactic Environment for the Heliosphere and Earth*. Dordrecht, The Netherlands: Springer, Astrophysics and Space Science Library, Vol. 338, 133–193.
- Genthon, C., Barnola, J. M., Raynaud, D., Lorius, C., Jouzel, J., Barkov, N. I., Korotkevich, Y. S., and Kotlyakov, V. M. (1987) Vostok ice core: Climatic response

- to CO₂ and orbital forcing changes over the last climatic cycle. *Nature*, Vol. 329, 414–418.
- Gore, D. B., Rhodes, E. J., Augustinus, P. C., Leishman, M. R., Colhoun, E. A., and Rees-Jones, J. (2001) Bunge Hills, East Antarctica: Ice free at the last glacial maximum. *Geology*, Vol. 29, 1103–1106.
- Gough, D. O. (1990) On possible origins of relatively short-term variations in the solar structure. *Philosophical Transactions of the Royal Society (London) A*, Vol. 330, 627–640.
- Gough, D. O. (2002) How is solar activity influencing the structure of the Sun? In A. Wilson, ed., *From Solar Min to Max: Half a Solar Cycle with SOHO*. Noordwijk, The Netherlands: ESA–European Space Agency, Proceedings of SOHO 11 Symposium (ESA SP-508), 577–592.
- Greene, A. M., Seager, R., and Broecker, W. S. (2002) Tropical snowline depression at the last glacial maximum: Comparison with proxy records using single-cell tropical climate model. *Journal of Geophysical Research*, Vol. 107, D84061, doi:10.1029/2001JD000670.
- Gupta, S. M., Fernandes, A. A., and Mohan, R. (1996) Tropical sea surface temperatures and Earth's orbital eccentricity cycles. *Geophysical Research Letters*, Vol. 22, 3159–3162.
- Hansen, J., Lacis, A., Ruedy, R., Sato, M., and Wilson, H. (1993) How sensitive is the world's climate? *National Geographic Research & Exploration*, Vol. 9, No. 2, 142–158.
- Hansen, J., Sato, M., Lacis, A., and Ruedy, R. (1997) The missing climate forcing. *Philosophical Transactions of the Royal Society (London) B*, Vol. 352, 231–240.
- Hansen, J., Sato, M., Kharecha, P., Russell, G., Lea, D. W., and Siddall, M. (2007) Climate change and trace gases. *Philosophical Transactions of the Royal Society*, Vol. 365, 1925–1954.
- Haywood, A. M., Valdes, P. J., and Peck, V. L. (2007) A permanent El Niño-like state during the Pliocene? *Paleocenography*, Vol. 22, PA1213, doi:10.1029/2006PA001323.
- Hewitt, C. D. and Mitchell, J. F. B. (1997) Radiative forcing and response of a GCM to ice age boundary conditions: Cloud feedback and climate sensitivity. *Climate Dynamics*, Vol. 13, 821–834.
- Hodgson, D. A., Verleyen, E., Squier, A. H., Sabbe, K., Keely, B. J., Saunders, K. M., and Vyverman, W. (2006) Interglacial environments of coastal east Antarctica: Comparison of MIS 1 (Holocene) and MIS 5e (Last Interglacial) lake sediment records. *Quaternary Science Reviews*, Vol. 25, 179–197.
- Holzhauser, H., Magny, M., and Zumbuhl, H. J. (2005) Glacier and lake-level variations in west-central Europe over the last 3500 years. *Holocene*, Vol. 15, 789–801.
- Holzhammer, S., Mangini, A., Spotl, C., and Mudelsee, M. (2004) Timing and progression of the Last Interglacial derived from a high alpine stalagmite. *Geophysical Research Letters*, Vol. 31, L07201, doi:10.1029/2003GL019112.
- Huybers, P. (2007) Glacial variability over the last two million years: An extended depth-derived age model, continuous obliquity pacing, and the Pleistocene progression. *Quaternary Science Reviews*, Vol. 26, 37–55.

- Huybers, P. and Molnar, P. (2007) Tropical cooling and the onset of North American glaciation. *Climate of the Past Discussion*, Vol. 3, 771–789.
- Huybers, P. and Wunsch, C. (2005) Obliquity pacing of the late Pleistocene glacial terminations. *Nature*, Vol. 434, 491–494.
- Ikehara, M., Kawamura, K., Ohkouchi, N., Kimoto, K., Murayama, M., Nakamura, T., Oba, T., and Taira, A. (1997) Alkenone sea surface temperature in Southern Ocean for the last two deglaciations. *Geophysical Research Letters*, Vol. 24, 679–682.
- IPCC. (2007) *Climate Change 2007: The Physical Science Basis* (Working Group I contribution to the UN IPCC Fourth Assessment Report available at <http://www.ipcc.ch>). Cambridge, UK, and New York, NY: Cambridge University Press.
- Ishiwatari, R., Houtatsu, M., and Okada, H. (2001) Alkenone-sea surface temperatures in the Japan Sea over the past 36 kyr: Warm temperatures at the last glacial maximum. *Organic Geochemistry*, Vol. 32, 57–67.
- Johnsen, S. J., Dahl-Jensen, D., Gundestrup, N., Steffensen, J. P., Clausen, H. B., Miller, H., Masson-Delmotte, V., Sveinbjornsdottir, A. E., and White, J. (2001) Possible role for dust or other northern forcing of ice-age carbon dioxide changes. *Journal of Quaternary Science*, Vol. 16, 299–307.
- Johnston, T. C. and Alley, R. B. (2006) Possible role for dust or other northern forcing of ice-age carbon dioxide changes. *Quaternary Science Reviews*, Vol. 25, 3198–3206.
- Joos, F. (2005) Radiative forcing and the ice core greenhouse gas record. *PAGES News*, Vol. 13, No. 3, 11–13.
- Joshi, M., Shine, K., Ponater, M., Stuber, N., Rausen, R., and Li, L. (2003) A comparison of climate response to different radiative forcings in three general circulation models: Towards an improved metric of climate change. *Climate Dynamics*, Vol. 20, 843–854.
- Kato, S., Loeb, N. G., Minnis, P., Francis, J. A., Charlock, T. P., Rutan, D. A., Clothiaux, E. E., and Sun-Mack, S. (2006) Seasonal and interannual variations of top-of-atmosphere irradiance and cloud cover over polar regions derived from CERES data set. *Geophysical Research Letters*, Vol. 33, L19804, doi:10.1029/2006GL026685.
- Kaufman, D. S. and 29 co-authors. (2004) Holocene thermal maximum in the western Arctic (0–180°W). *Quaternary Science Reviews*, Vol. 23, 529–560.
- Kawamura, K. and 17 co-authors. (2007) Northern Hemisphere forcing of climate cycles in Antarctica over the past 360,000 years. *Nature*, Vol. 448, 912–916.
- Khodri, M., Leclainche, Y., Ramstein, G., Braconnot, P., Marti, O., and Cortijo, E. (2001) Simulating the amplification of orbital forcing by ocean feedbacks in the last glaciation. *Nature*, Vol. 410, 570–574.
- Khodri, M., Ramstein, G., de Noblet-Ducoudré, N., and Kageyama, M. (2003) Sensitivity of the northern extratropics hydrological cycle to the changing insolation forcing at 126 and 115 ky BP. *Climate Dynamics*, Vol. 21, 273–287.
- Kim, J.-H., Meggers, H., Rimbu, N., Lohmann, G., Freudenthal, T., Muller, P. J., and Schneider, R. R. (2007) Impacts of the North Atlantic gyre circulation on Holocene climate off northwest Africa. *Geology*, Vol. 35, 387–390.

- Kobashi, T., Severinghaus, J. P., Brook, E. J., Barnola, J.-M., and Grachev, A. M. (2007) Precise timing and characterization of abrupt climate change 8200 years ago from air trapped in polar ice. *Quaternary Science Reviews*, Vol. 26, 1212–1222.
- Kubatzki, C., Claussen, M., Calov, R., and Ganopolski, A. (2006) Sensitivity of the last glacial inception to initial and surface conditions. *Climate Dynamics*, Vol. 27, 333–344.
- Kukla, G. and Gavin, J. (2005) Did glacials start with global warming? *Quaternary Science Reviews*, Vol. 24, 1547–1557.
- Lal, D., Jull, A. J. T., Pollard, D., and Vacher, L. (2005) Evidence for large century time-scale changes in solar activity in the past 32 kyr, based on in-situ cosmogenic ^{14}C in ice at Summit, Greenland. *Earth and Planetary Science Letters*, Vol. 234, 335–349.
- Laskar, J., Joutel, F., and Boudin, F. (1993) Orbital, precessional, and insolation quantities for the Earth from -20 Myr to +10 Myr. *Astronomy and Astrophysics*, Vol. 270, 522–533.
- Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correira, A. C. M., and Levrard, B. (2004) A long-term numerical solution for the insolation quantities of the Earth. *Astronomy and Astrophysics*, Vol. 428, 261–285.
- Lea, D. W. (2004) The 100000-yr cycle in tropical SST, greenhouse forcing, and climate sensitivity. *Journal of Climate*, Vol. 17, 2170–2179.
- Lea, D. W., Pak, D. K., and Spero, H. J. (2000) Climate impact of late Quaternary equatorial Pacific sea surface temperature variations. *Science*, Vol. 289, 1719–1724.
- Leduc, G., Vidal, L., Tachikawa, K., Rostek, F., Sonzogni, C., Beaufort, L., and Bard, E. (2007) Moisture transport across Central America as a positive feedback on abrupt climatic changes. *Nature*, Vol. 445, 908–911.
- Lisiecki, L. E. and Raymo, M. E. (2007) Plio-Pleistocene climate evolution: Trends and transitions in glacial cycle dynamics. *Quaternary Science Reviews*, Vol. 26, 56–69.
- Liu, Z. and Herbert, T. D. (2004) High-latitude influence on the eastern equatorial Pacific climate in the early Pleistocene epoch. *Nature*, Vol. 427, 720–723.
- Lorenz, S. J., Kim, J.-H., Rimbu, N., Schneider, R. R., and Lohmann, G. (2006) Orbitally driven insolation forcing on Holocene climate trends: Evidence from alkenone data and climate modelling. *Paleoenography*, Vol. 21, PA1002, doi:10.1029/2005PA001152.
- Lorius, C., Jouzel, J., Raynaud, D., Hansen, J., and Le Treut, H. (1990) The ice-core record: Climate sensitivity and future greenhouse warming. *Nature*, Vol. 347, 139–145.
- Loulergue, L., Parrenin, F., Blunier, T., Barnola, J.-M., Spahni, R., Schlit, A., Raisbeck, G., and Chappellaz, J. (2007) New constraints on the gas age–ice age difference along the EPICA ice cores, 0 to 50 kyr. *Climate of the Past Discussion*, Vol. 3, 435–467.
- Loutre, M. F. and Berger, A. (2000) No glacial–interglacial cycle in the ice volume simulated under a constant astronomical forcing and a variable CO_2 . *Geophysical Research Letters*, Vol. 27, 783–786.

- Loutre, M. F., Berger, A., Bretagnon, P., and Blanc, P. L. (1992) Astronomical frequencies for climate research at the decadal to century time scale. *Climate Dynamics*, Vol. 7, 181–194.
- Loutre, M. F., Paillard, D., Vimeux, F., and Cortijo, E. (2004) Does mean annual insolation have the potential to change the climate? *Earth and Planetary Science Letters*, Vol. 221, 1–14.
- Maasch, K. A., Mayweski, P. A., Rohling, E. J., Stager, J. C., Karlen, W., Meeker, L. D., and Meyerson, E. A. (2005) A 2000-year context for modern climate change. *Geografiska Annaler*, Vol. 87, 7–15.
- Magny, M. and 11 co-authors. (2007) Holocene climate change in the central Mediterranean as recorded by lake-level fluctuations at lake Accesa (Tuscany, Italy). *Quaternary Science Reviews*, Vol. 26, 1736–1758.
- Marchal, O. and 18 co-authors. (2002) Apparent long-term cooling of the sea surface in the northeast Atlantic and Mediterranean during the Holocene. *Quaternary Science Reviews*, Vol. 21, 455–483.
- Martinson, D. G. and Pitman III, W. C. (2007) The Arctic as a trigger for glacial terminations. *Climatic Change*, Vol. 80, 253–263.
- Masson, V. and 13 co-authors. (2000) Holocene climate variability in Antarctica based on 11 ice-core isotopic records. *Quaternary Research*, Vol. 54, 348–358.
- Masson-Delmotte, V., Dreyfus, G., Braconnot, P., Johnsen, S., Jouzel, J., Kageyama, M., Landais, A., Loutre, M.-F., Nouet, J., Parrenin, F., Raynaud, D., Stenni, B., and Tüentler, E. (2006) Past temperature reconstructions from deep ice cores: Relevance for future climate change. *Climate of the Past*, Vol. 2, 145–165.
- Masson-Delmotte, V., Stenni, B., and Jouzel, J. (2004) Common millennial-scale variability of Antarctic and Southern Ocean temperatures during the past 5000 years reconstructed from the EPICA Dome C ice core. *Holocene*, Vol. 14, 145–151.
- Mayewski, P. A. and 15 co-authors. (2004) Holocene climate variability. *Quaternary Research*, Vol. 62, 243–255.
- Milkov, A. V. (2004) Global estimates of hydrate-bound gas in marine sediments: How much is really out there? *Earth-Science Reviews*, Vol. 66, 183–197.
- Monnin, E., Indermuhle, A., Dallenbach, A., Flückiger, J., Stauffer, B., Stocker, T. F., Raynaud, D., and Barnola, J.-M. (2001) Atmospheric CO₂ concentrations over the last glacial termination. *Science*, Vol. 291, 112–114.
- Mudelsee, M. (2001) The phase relations among atmospheric CO₂ content, temperature and global ice volume over the past 420 ka. *Quaternary Science Reviews*, Vol. 20, 583–589.
- Müller, H.-R., Frisch, P. C., Florinski, V., and Zank, G. P. (2006) Heliospheric response to different possible interstellar environments. *Astrophysical Journal*, Vol. 647, 1491–1505.
- Norgaard-Pedersen, N., Spielhagen, R. F., Erlenkeuser, H., Grootes, P. M., Heinemeier, J., and Knies, J. (2003) Arctic ocean during the Last Glacial Maximum: Atlantic and polar domains of surface water mass distribution and ice cover. *Paleoceanography*, Vol. 18, 1063, doi:10.1029/2002PA000781.

- Overpeck, J., Rind, D., Lacy, A., and Healy, R. (1996) Possible role of dust-induced regional warming in abrupt climate change during the last glacial period. *Nature*, Vol. 384, 447–449.
- Pahnke, K. and Sachs, J. P. (2006) Sea surface temperatures of southern midlatitudes 0–160 kyr B.P. *Paleoceanography*, Vol. 21, PA2003, doi:10.1029/2005PA001191.
- Peacock, S., Lane, E., and Restrepo, J. M. (2006) A possible sequence of events for the generalized glacial–interglacial cycle. *Global Biogeochemical Cycles*, Vol. 20, GB2010, doi:10.1029/2005GB002448.
- Peltier, W. R. and Solheim, L. P. (2004) The climate of the Earth at Last Glacial Maximum: Statistical equilibrium state and a mode of internal variability. *Quaternary Science Reviews*, Vol. 23, 335–357.
- Pepin, L., Raynaud, D., Barnola, J. M., and Loutre, M. F. (2001) Hemispheric roles of climate forcings during glacial–interglacial transitions as deduced from the Vostok record and LLN-2D model experiments. *Journal of Geophysical Research*, Vol. 106, 31885–31892.
- Petit, J. R. and 18 co-authors. (1999) Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica. *Nature*, Vol. 399, 429–436.
- Posmentier, E. S. (1994) Response of an ocean-atmosphere climate model to Milankovic forcing. *Nonlinear Processes in Geophysics*, Vol. 1, 26–30.
- Priem, H. N. A. (1997) CO₂ and climate: A geologist's view. *Space Science Reviews*, Vol. 81, 173–198.
- Ravelo, A. C., Andreasen, D. H., Lyle, M., Lyle, A. O., and Wara, M. W. (2004) Regional climate shifts caused by gradual cooling in the Pliocene epoch. *Nature*, Vol. 429, 263–267.
- Raymo, M. E. and Nisancioglu, K. (2003) The 41kyr world: Milankovitch's other unresolved mystery. *Paleoceanography*, Vol. 18, doi:10.1029/2002PA000791.
- Richardson, M. I. and Mischna, M. A. (2005) Long-term evolution of transient liquid water on Mars. *Journal of Geophysical Research*, Vol. 110, E03003, doi:10.1029/2004JE002367.
- Risebrobakken, B., Dokken, T., Ottera, O. H., Jansen, E., Gao, Y., and Drange, H. (2007) Inception of the Northern European ice sheet due to contrasting ocean and insolation forcing. *Quaternary Research*, Vol. 67, 128–135.
- Roe, G. (2006) In defense of Milankovitch. *Geophysical Research Letters*, Vol. 33, L24703, doi:10.1029/2006GL027817.
- Ruddiman, W. F. and Raymo, M. E. (2003) A methane-based time scale for Vostok ice. *Quaternary Science Reviews*, Vol. 22, 141–155.
- Sachs, J. P. (2007) Cooling of Northwest Atlantic slope waters during the Holocene. *Geophysical Research Letters*, Vol. 33, L03609, doi:10.1029/2006GL028495.
- Saltzman, B., Maasch, K. A., and Verbitsky, M. Ya. (1993) Possible effects of anthropogenically-increased CO₂ on the dynamics of climate: Implications for ice age cycles. *Geophysical Research Letters*, Vol. 20, 1051–1054.
- Schaefer, H., Whiticar, M. J., Brook, E. J., Petrenko, V. V., Ferretti, D. F., and Severinghaus, J. P. (2006) Ice record of $\delta^{13}\text{C}$ for atmospheric CH₄ across the Younger Dryas-Preboreal transition. *Science*, Vol. 313, 1109–1112.

- Scherer, K. and 13 co-authors. (2006) Interstellar-terrestrial relations: Variable cosmic environments, the dynamic heliosphere, and their imprints on terrestrial archives and climate. *Space Science Reviews*, Vol. 127, 327–465.
- Schneider von Deimling, T., Ganopolski, A., Held, H., and Rahmstorf, S. (2006a) How cold was the last glacial maximum? *Geophysical Research Letters*, Vol. 33, L14709, doi:10.1029/2006GL026484.
- Schneider von Deimling, T., Held, H., Ganopolski, A., and Rahmstorf, S. (2006b) Climate sensitivity estimated from ensemble simulations of glacial climate. *Climate Dynamics*, Vol. 27, 149–163.
- Schrag, D. P., Hampt, G., and Murray, D. W. (1996) Pore fluid constraints on the temperature and oxygen isotopic composition of the glacial ocean. *Science*, Vol. 272, 1930–1932.
- Seki, O., Kawamura, K., Ikehara, M., Nakatsuka, T., and Oba, T. (2004) Variation of alkenone sea surface temperature in the Sea of Okhotsk over the last 85 kyrs. *Organic Geochemistry*, Vol. 35, 347–354.
- Shackleton, N. J. (2000) The 100,000-year ice-age cycle identified and found to lag temperature, carbon dioxide, and orbital eccentricity. *Science*, Vol. 289, 1897–1902.
- Sharma, M. (2002) Variations in solar magnetic activity during the last 200000 years: Is there a Sun-climate connection? *Earth and Planetary Science Letters*, Vol. 199, 459–472.
- Shell, K. M., Frouin, R., Nakamoto, S., and Sommerville, R. C. J. (2003) Atmospheric response to solar radiation absorbed by phytoplankton. *Journal of Geophysical Research*, Vol. 108, D15, doi:10.1029/2003JD003440.
- Siegenthaler, U., Stocker, T. F., Monnin, E., Luthi, D., Schwander, J., Stauffer, B., Raynaud, D., Barnola, J.-M., Fischer, H., Masson-Delmotte, V., and Jouzel, J. (2005) Stable carbon cycle–climate relationship during the late Pleistocene. *Science*, Vol. 310, 1313–1317.
- Skinner, L. C. (2006) Glacial–interglacial atmospheric CO₂ change: A simple “hypso-metric effect” on deep-ocean carbon sequestration? *Climate of the Past Discussions*, Vol. 2, 711–743.
- Smith, J. A., Bentley, M. J., Hodgson, D. A., Roberts, S. J., Leng, M. J., Lloyd, J. M., Barrett, M. S., Bryant, C., and Sugden, D. E. (2007) Oceanic and atmospheric forcing of early Holocene ice shelf retreat, George VI Ice Shelf, Antarctic Peninsula. *Quaternary Science Reviews*, Vol. 26, 500–516.
- Soon, W. W.-H. (2005) Variable solar irradiance as a plausible agent for multidecadal variations in the Arctic-wide surface air temperature record of the past 130 years. *Geophysical Research Letters*, Vol. 32, L16712, doi:10.1029/2005GL023429.
- Soon, W., Baliunas, S., Idso, S. B., Kondratyev, K. Ya., and Posmentier, E. S. (2001) Modeling climatic effects of anthropogenic carbon dioxide emissions: Unknowns and uncertainties. *Climate Research*, Vol. 18, 259–275.
- Spahni, R., Chappellaz, J., Stocker, T. F., Loulergue, L., Hausammann, G., Kawamura, K., Flückiger, J., Schwander, J., Raynaud, D., Masson-Delmotte, V., and Jouzel, J. (2005) Atmospheric methane and nitrous oxide of the late Pleistocene from Antarctic ice cores. *Science*, Vol. 310, 1317–1321.

- Spielhagen, R. F. and 14 co-authors. (1997) Arctic ocean evidence for late Quaternary initiation of northern Eurasian ice sheets. *Geology*, Vol. 25, 783–786.
- Stenni, B., Masson-Delmotte, V., Johnsen, S., Jouzel, J., Longinelli, A., Monnin, E., Rothlisberger, R., and Selmo, E. (2001) An oceanic cold reversal during the last deglaciation. *Science*, Vol. 293, 2074–2077.
- Stott, L., Cannariato, K., Thunell, R., Haug, G. H., Koutavas, A., and Lund, S. (2004) Decline of surface temperature and salinity in the western tropical Pacific ocean in the Holocene epoch. *Nature*, Vol. 431, 56–59.
- Suggate, R. P. and Almond, P. C. (2005) The Last Glacial Maximum (LGM) in western South Island, New Zealand: Implications for the global LGM and MIS 2. *Quaternary Science Reviews*, Vol. 24, 1923–1940.
- Sutherland, R., Kim, K., Zondervan, A., and McSaveney, M. (2007) Orbital forcing of mid-latitude Southern Hemisphere glaciation since 100 ka inferred from cosmogenic nuclide ages of moraine boulders from the Cascase Plateau, southwest New Zealand. *Geological Society of America Bulletin*, Vol. 119, 443–451.
- Svendsen, J. I. and 13 co-authors. (1999) Maximum extent of the Eurasian ice sheets in the Barents and Kara Sea region during the Weichselian. *Boreas*, Vol. 28, 234–242.
- Turck-Chieze, S. and 39 co-authors. (2005) The magnetism of the solar interior for a complete MHD solar vision. In F. Favata, J. Sanz-Forcada, and A. Gimenez, eds., *Trends in Space Science and Cosmic Vision 2020*. Noordwijk, The Netherlands: ESA–European Space Agency, Proceedings of 2005 ESLAB Symposium (ESA SP-588), 193–202.
- Tziperman, E., Raymo, M. E., Huybers, P., and Wunsch, C. (2006) Consequences of pacing the Pleistocene 100 kyr ice ages by nonlinear phase locking to Milankovitch forcing. *Paleocenography*, Vol. 21, PA4206, doi:10.1029/2005PA001241.
- Vallina, S. M. and Simo, R. (2007) Strong relationship between DMS and the solar radiation dose over the global surface ocean. *Science*, Vol. 315, 506–508.
- Vandergoes, M. J., Newnham, R. M., Preusser, F., Hendy, C. H., Lowell, T. V., Fitzsimons, S. J., Hogg, A. G., Kasper, H. U., and Schluchter, C. (2005) Regional insolation forcing of late Quaternary climate change in the Southern Hemisphere. *Nature*, Vol. 436, 242–245.
- Vettoretti, G. and Peltier, W. R. (2003) Post-Eemian glacial inception. Part II: Elements of a cryospheric moisture pump. *Journal of Climate*, Vol. 16, 912–927.
- Vettoretti, G. and Peltier, W. R. (2004) Sensitivity of glacial inception to orbital and greenhouse gas climate forcing. *Quaternary Science Reviews*, Vol. 23, 499–519.
- Vimeux, F., Cuffey, K. M., and Jouzel, J. (2002) New insights into Southern Hemisphere temperature changes from Vostok ice cores using deuterium excess correction. *Earth and Planetary Science Letters*, Vol. 203, 829–843.
- Visser, K., Thunell, R., and Stott, L. (2003) Magnitude and timing of temperature change in the Indo-Pacific warm pool during deglaciation. *Nature*, Vol. 421, 152–155.
- Weaver, A. J., Eby, M., Fanning, A., and Wiebe, E. C. (1998) Simulated influence of carbon dioxide, orbital forcing and ice sheets on the climate of the last glacial maximum. *Nature*, Vol. 394, 847–853.

- Weldeab, S., Lea, D. W., Schneider, R. R., and Andersen, N. (2007) 155,000 years of west African monsoon and ocean thermal evolution. *Science*, Vol. 316, 1303–1307.
- Yoshimori, M., Weaver, A. J., Marshall, S. J., and Clarke, G. K. C. (2001) Glacial termination: Sensitivity to orbital and CO₂ forcing in a coupled climate system model. *Climate Dynamics*, Vol. 17, 571–588.
- Yung, Y. L., Lee, T., Wang, C.-H., and Shieh, Y.-T. (1996) Dust: A diagnostic of the hydrologic cycle during the Last Glacial Maximum. *Science*, Vol. 271, 962–963.
- Zhang, Y., Rossow, W. B., Lacis, A. A., Oinas, V., and Mishchenko, M. I. (2004) Calculation of radiative fluxes from the surface to top of atmosphere based on ISCCP and other global data sets: Refinements of the radiative transfer model and the input data. *Journal of Geophysical Research*, Vol. 109, D19105, doi:10.1029/2003JD004457.